

Reference list

- Albers,S.V., Elferink,M.G., Charlebois,R.L., Sensen,C.W., Driessen,A.J., and Konings,W.N. (1999) Glucose transport in the extremely thermoacidophilic *Sulfolobus solfataricus* involves a high-affinity membrane-integrated binding protein. *J Bacteriol* **181**: 4285-4291.
- Albers,S.V., Jonuscheit,M., Dinkelaker,S., Ulrich,T., Kletzin,A., Tampe,R. *et al.* (2006) Production of recombinant and tagged proteins in the hyperthermophilic archaeon *Sulfolobus solfataricus*. *Appl Environ Microbiol* **72**: 102-111.
- Albers,S.V., Koning,S.M., Konings,W.N., and Driessen,A.J. (2004) Insights into ABC transport in archaea. *J Bioenerg Biomembr* **36**: 5-15.
- Albers,S.V., Van de Vossenberg, Driessen,A.J., and Konings,W.N. (2001) Bioenergetics and solute uptake under extreme conditions. *Extremophiles* **5**: 285-294.
- Albers,S.-V., and Driessen,A.J.M. (2008) Conditions for Gene Disruption by Homologous Recombination of Exogenous DNA into the *Sulfolobus solfataricus* Genome. *Archaea* **2**: 145-149.
- Allers,T., and Mevarech,M. (2005) Archaeal genetics - the third way. *Nat Rev Genet* **6**: 58-73.
- Aravind,L., Anantharaman,V., Balaji,S., Babu,M.M., and Iyer,L.M. (2005) The many faces of the helix-turn-helix domain: transcription regulation and beyond. *FEMS Microbiol Rev* **29**: 231-262.
- Aravind,L., and Koonin,E.V. (1999) DNA-binding proteins and evolution of transcription regulation in the archaea. *Nucleic Acids Res* **27**: 4658-4670.
- Baliga,N.S., and Dassarma,S. (1999) Saturation mutagenesis of the TATA box and upstream activator sequence in the haloarchaeal *bop* gene promoter. *J Bacteriol* **181**: 2513-2518.
- Baliga,N.S., Goo,Y.A., Ng,W.V., Hood,L., Daniels,C.J., and Dassarma,S. (2000) Is gene expression in *Halobacterium* NRC-1 regulated by multiple TBP and TFB transcription factors? *Mol Microbiol* **36**: 1184-1185.
- Baliga,N.S., Kennedy,S.P., Ng,W.V., Hood,L., and Dassarma,S. (2001) Genomic and genetic dissection of an archaeal regulon. *Proc Natl Acad Sci U S A* **98**: 2521-2525.
- Banerjee,S., Chalissery,J., Bandey,I., and Sen,R. (2006) Rho-dependent transcription termination: more questions than answers. *J Microbiol* **44**: 11-22.
- Barns,S.M., Delwiche,C.F., Palmer,J.D., and Pace,N.R. (1996) Perspectives on archaeal diversity, thermophily and monophyly from environmental rRNA sequences. *Proc Natl Acad Sci U S A* **93**: 9188-9193.
- Barry,E.R., and Bell,S.D. (2006) DNA replication in the archaea. *Microbiol Mol Biol Rev* **70**: 876-887.
- Barthelme,D., Scheele,U., Dinkelaker,S., Janoschka,A., Macmillan,F., Albers,S.V. *et al.* (2007) Structural organization of essential iron-sulfur clusters in the evolutionarily highly conserved ATP-binding cassette protein ABCE1. *J Biol Chem* **282**: 14598-14607.

Bell,S.D., Botting,C.H., Wardleworth,B.N., Jackson,S.P., and White,M.F. (2002) The interaction of Alba, a conserved archaeal chromatin protein, with Sir2 and its regulation by acetylation. *Science* **296**: 148-151.

Bell,S.D., Brinkman,A.B., Van der,O.J., and Jackson,S.P. (2001) The archaeal TFIIIEalpha homologue facilitates transcription initiation by enhancing TATA-box recognition. *EMBO Rep* **2**: 133-138.

Bell,S.D., Cairns,S.S., Robson,R.L., and Jackson,S.P. (1999a) Transcriptional regulation of an archaeal operon in vivo and in vitro. *Mol Cell* **4**: 971-982.

Bell,S.D., and Jackson,S.P. (2000b) Mechanism of autoregulation by an archaeal transcriptional repressor. *J Biol Chem* **275**: 31624-31629.

Bell,S.D., and Jackson,S.P. (2000a) The role of transcription factor B in transcription initiation and promoter clearance in the archaeon *Sulfolobus acidocaldarius*. *J Biol Chem* **275**: 12934-12940.

Bell,S.D., and Jackson,S.P. (2001) Mechanism and regulation of transcription in archaea. *Curr Opin Microbiol* **4**: 208-213.

Bell,S.D., Jaxel,C., Nadal,M., Kosa,P.F., and Jackson,S.P. (1998) Temperature, template topology, and factor requirements of archaeal transcription. *Proc Natl Acad Sci U S A* **95**: 15218-15222.

Bell,S.D., Kosa,P.L., Sigler,P.B., and Jackson,S.P. (1999b) Orientation of the transcription preinitiation complex in archaea. *Proc Natl Acad Sci U S A* **96**: 13662-13667.

Berkner,S., Grogan,D., Albers,S.V., and Lipps,G. (2007) Small multicopy, non-integrative shuttle vectors based on the plasmid pRN1 for *Sulfolobus acidocaldarius* and *Sulfolobus solfataricus*, model organisms of the (cren-)archaea. *Nucleic Acids Res* **35**: e88.

Best,A.A., and Olsen,G.J. (2001) Similar subunit architecture of archaeal and eukaryal RNA polymerases. *FEMS Microbiol Lett* **195**: 85-90.

Bohm,A., Diez,J., Diederichs,K., Welte,W., and Boos,W. (2002) Structural model of MalK, the ABC subunit of the maltose transporter of *Escherichia coli*: implications for mal gene regulation, inducer exclusion, and subunit assembly. *J Biol Chem* **277**: 3708-3717.

Boos,W., and Bohm,A. (2000) Learning new tricks from an old dog: MalT of the *Escherichia coli* maltose system is part of a complex regulatory network. *Trends Genet* **16**: 404-409.

Boos,W., and Shuman,H. (1998) Maltose/maltodextrin system of *Escherichia coli*: transport, metabolism, and regulation. *Microbiol Mol Biol Rev* **62**: 204-229.

Brinkman,A.B., Bell,S.D., Lebbink,R.J., de Vos,W.M., and Van der,O.J. (2002) The *Sulfolobus solfataricus* Lrp-like protein LysM regulates lysine biosynthesis in response to lysine availability. *J Biol Chem* **277**: 29537-29549.

Brinkman,A.B., Dahlke,I., Tuininga,J.E., Lammers,T., Dumay,V., de,H.E. *et al.* (2000) An Lrp-like transcriptional regulator from the archaeon *Pyrococcus furiosus* is negatively autoregulated. *J Biol Chem* **275**: 38160-38169.

- Brinkman,A.B., Ettema,T.J., de Vos,W.M., and Van der,O.J. (2003) The Lrp family of transcriptional regulators. *Mol Microbiol* **48**: 287-294.
- Brochier,C., Gribaldo,S., Zivanovic,Y., Confalonieri,F., and Forterre,P. (2005) Nanoarchaea: representatives of a novel archaeal phylum or a fast-evolving euryarchaeal lineage related to Thermococcales? *Genome Biol* **6**: R42.
- Brochier-Armanet,C., Boussau,B., Gribaldo,S., and Forterre,P. (2008) Mesophilic Crenarchaeota: proposal for a third archaeal phylum, the Thaumarchaeota. *Nat Rev Microbiol* **6**: 245-252.
- Brock,T.D., Brock,K.M., Belly,R.T., and Weiss,R.L. (1972) Sulfolobus: a new genus of sulfur-oxidizing bacteria living at low pH and high temperature. *Arch Mikrobiol* **84**: 54-68.
- Brouns,S.J., Walther,J., Snijders,A.P., van de Werken,H.J., Willems,H.L., Worm,P. *et al.* (2006) Identification of the missing links in prokaryotic pentose oxidation pathways: evidence for enzyme recruitment. *J Biol Chem* **281**: 27378-27388.
- Bukau,B., Ehrmann,M., and Boos,W. (1986) Osmoregulation of the maltose regulon in *Escherichia coli*. *J Bacteriol* **166**: 884-891.
- Burke,T.W., Willy,P.J., Kutach,A.K., Butler,J.E., and Kadonaga,J.T. (1998) The DPE, a conserved downstream core promoter element that is functionally analogous to the TATA box. *Cold Spring Harb Symp Quant Biol* **63**: 75-82.
- Chen,J., Lu,G., Lin,J., Davidson,A.L., and Quiococho,F.A. (2003) A tweezers-like motion of the ATP-binding cassette dimer in an ABC transport cycle. *Mol Cell* **12**: 651-661.
- Chen,L., Brugger,K., Skovgaard,M., Redder,P., She,Q., Torarinsson,E. *et al.* (2005) The genome of *Sulfolobus acidocaldarius*, a model organism of the Crenarchaeota. *J Bacteriol* **187**: 4992-4999.
- Cohen-Kupiec,R., Blank,C., and Leigh,J.A. (1997) Transcriptional regulation in Archaea: in vivo demonstration of a repressor binding site in a methanogen. *Proc Natl Acad Sci U S A* **94**: 1316-1320.
- Cohen-Kupiec,R., Marx,C.J., and Leigh,J.A. (1999) Function and regulation of *glnA* in the methanogenic archaeon *Methanococcus maripaludis*. *J Bacteriol* **181**: 256-261.
- Cohen-Kupiec,R., Zilberstein,A., and Gurevitz,M. (1995) Characterization of cis elements that regulate the expression of *glnA* in *Synechococcus* sp. strain PCC 7942. *J Bacteriol* **177**: 2222-2226.
- Coleman,R.A., and Pugh,B.F. (1997) Slow dimer dissociation of the TATA binding protein dictates the kinetics of DNA binding. *Proc Natl Acad Sci U S A* **94**: 7221-7226.
- Coleman,R.A., Taggart,A.K., Burma,S., Chicca,J.J., and Pugh,B.F. (1999) TFIIA regulates TBP and TFIID dimers. *Mol Cell* **4**: 451-457.
- Collier,D.N., Hager,P.W., and Phibbs,P.V., Jr. (1996) Catabolite repression control in the Pseudomonads. *Res Microbiol* **147**: 551-561.
- Cubonova,L., Sandman,K., Hallam,S.J., Delong,E.F., and Reeve,J.N. (2005) Histones in crenarchaea. *J Bacteriol* **187**: 5482-5485.

- Dahlke,I., and Thomm,M. (2002) A Pyrococcus homolog of the leucine-responsive regulatory protein, LrpA, inhibits transcription by abrogating RNA polymerase recruitment. *Nucleic Acids Res* **30**: 701-710.
- Darcy,T.J., Hausner,W., Awery,D.E., Edwards,A.M., Thomm,M., and Reeve,J.N. (1999) Methanobacterium thermoautotrophicum RNA polymerase and transcription in vitro. *J Bacteriol* **181**: 4424-4429.
- Decker,B.S., O'Brien,R., Fleming,P.J., Geiger,J.H., Jackson,S.P., and Sigler,P.B. (1996) The crystal structure of a hyperthermophilic archaeal TATA-box binding protein. *J Mol Biol* **264**: 1072-1084.
- den Hengst,C.D., van Hijum,S.A., Geurts,J.M., Nauta,A., Kok,J., and Kuipers,O.P. (2005) The Lactococcus lactis CodY regulon: identification of a conserved cis-regulatory element. *J Biol Chem* **280**: 34332-34342.
- Diederichs,K., Diez,J., Greller,G., Muller,C., Breed,J., Schnell,C. *et al.* (2000) Crystal structure of Malk, the ATPase subunit of the trehalose/maltose ABC transporter of the archaeon Thermococcus litoralis. *EMBO J* **19**: 5951-5961.
- Egorova,K., and Antranikian,G. (2005) Industrial relevance of thermophilic Archaea. *Curr Opin Microbiol* **8**: 649-655.
- Elferink,M.G., Albers,S.V., Konings,W.N., and Driessen,A.J. (2001) Sugar transport in Sulfolobus solfataricus is mediated by two families of binding protein-dependent ABC transporters. *Mol Microbiol* **39**: 1494-1503.
- Englert,C., Wanner,G., and Pfeifer,F. (1992) Functional analysis of the gas vesicle gene cluster of the halophilic archaeon Haloferax mediterranei defines the vac-region boundary and suggests a regulatory role for the gvpD gene or its product. *Mol Microbiol* **6**: 3543-3550.
- Enoru-Eta,J., Gigot,D., Glansdorff,N., and Charlier,D. (2002) High resolution contact probing of the Lrp-like DNA-binding protein Ss-Lrp from the hyperthermoacidophilic crenarchaeote Sulfolobus solfataricus P2. *Mol Microbiol* **45**: 1541-1555.
- Enoru-Eta,J., Gigot,D., Thia-Toong,T.L., Glansdorff,N., and Charlier,D. (2000) Purification and characterization of Sa-Lrp, a DNA-binding protein from the extreme thermoacidophilic archaeon Sulfolobus acidocaldarius homologous to the bacterial global transcriptional regulator Lrp. *J Bacteriol* **182**: 3661-3672.
- Ettema,T.J., Brinkman,A.B., Lamers,P.P., Kornet,N.G., de Vos,W.M., and Van der,O.J. (2006) Molecular characterization of a conserved archaeal copper resistance (cop) gene cluster and its copper-responsive regulator in Sulfolobus solfataricus P2. *Microbiology* **152**: 1969-1979.
- Fish,R.N., and Kane,C.M. (2002) Promoting elongation with transcript cleavage stimulatory factors. *Biochim Biophys Acta* **1577**: 287-307.
- Fox,G.E., Magrum,L.J., Balch,W.E., Wolfe,R.S., and Woese,C.R. (1977) Classification of methanogenic bacteria by 16S ribosomal RNA characterization. *Proc Natl Acad Sci U S A* **74**: 4537-4541.
- Goede,B., Naji,S., von,K.O., Ilg,K., and Thomm,M. (2006) Protein-protein interactions in the archaeal transcriptional machinery: binding studies of isolated RNA polymerase subunits and transcription factors. *J Biol Chem* **281**: 30581-30592.

- Grabowski,B., and Kelman,Z. (2003) Archeal DNA replication: eukaryal proteins in a bacterial context. *Annu Rev Microbiol* **57**: 487-516.
- Gregor,D., and Pfeifer,F. (2001) Use of a halobacterial bgaH reporter gene to analyse the regulation of gene expression in halophilic archaea. *Microbiology* **147**: 1745-1754.
- Grogan,D.W. (1989) Phenotypic characterization of the archaeobacterial genus Sulfolobus: comparison of five wild-type strains. *J Bacteriol* **171**: 6710-6719.
- Grogan,D.W., and Hansen,J.E. (2003) Molecular characteristics of spontaneous deletions in the hyperthermophilic archaeon Sulfolobus acidocaldarius. *J Bacteriol* **185**: 1266-1272.
- Gropp,F., and Betlach,M.C. (1994) The bat gene of Halobacterium halobium encodes a trans-acting oxygen inducibility factor. *Proc Natl Acad Sci U S A* **91**: 5475-5479.
- Hain,J., Reiter,W.D., Hudepohl,U., and Zillig,W. (1992) Elements of an archaeal promoter defined by mutational analysis. *Nucleic Acids Res* **20**: 5423-5428.
- Hallam,S.J., Konstantinidis,K.T., Putnam,N., Schleper,C., Watanabe,Y., Sugahara,J. *et al.* (2006) Genomic analysis of the uncultivated marine crenarchaeote Cenarchaeum symbiosum. *Proc Natl Acad Sci U S A* **103**: 18296-18301.
- Hampsey,M., and Reinberg,D. (1999) RNA polymerase II as a control panel for multiple coactivator complexes. *Curr Opin Genet Dev* **9**: 132-139.
- Haseltine,C., Montalvo-Rodriguez,R., Bini,E., Carl,A., and Blum,P. (1999a) Coordinate transcriptional control in the hyperthermophilic archaeon Sulfolobus solfataricus. *J Bacteriol* **181**: 3920-3927.
- Haseltine,C., Montalvo-Rodriguez,R., Carl,A., Bini,E., and Blum,P. (1999b) Extragenic pleiotropic mutations that repress glycosyl hydrolase expression in the hyperthermophilic archaeon Sulfolobus solfataricus. *Genetics* **152**: 1353-1361.
- Haseltine,C., Rolfmeier,M., and Blum,P. (1996) The glucose effect and regulation of alpha-amylase synthesis in the hyperthermophilic archaeon Sulfolobus solfataricus. *J Bacteriol* **178**: 945-950.
- Hausner,W., Frey,G., and Thomm,M. (1991) Control regions of an archaeal gene. A TATA box and an initiator element promote cell-free transcription of the tRNA(Val) gene of Methanococcus vannielii. *J Mol Biol* **222**: 495-508.
- Hausner,W., Lange,U., and Musfeldt,M. (2000) Transcription factor S, a cleavage induction factor of the archaeal RNA polymerase. *J Biol Chem* **275**: 12393-12399.
- Hausner,W., and Thomm,M. (2001) Events during initiation of archaeal transcription: open complex formation and DNA-protein interactions. *J Bacteriol* **183**: 3025-3031.
- Hausner,W., Wettach,J., Hethke,C., and Thomm,M. (1996) Two transcription factors related with the eucaryal transcription factors TATA-binding protein and transcription factor IIB direct promoter recognition by an archaeal RNA polymerase. *J Biol Chem* **271**: 30144-30148.

- Hoang,V., Bini,E., Dixit,V., Drozda,M., and Blum,P. (2004) The role of cis-acting sequences governing catabolite repression control of lacS expression in the archaeon *Sulfolobus solfataricus*. *Genetics* **167**: 1563-1572.
- Hochheimer,A., Hedderich,R., and Thauer,R.K. (1999) The DNA binding protein Tfx from *Methanobacterium thermoautotrophicum*: structure, DNA binding properties and transcriptional regulation. *Mol Microbiol* **31**: 641-650.
- Hofacker,A., Schmitz,K.M., Cichonczyk,A., Sartorius-Neef,S., and Pfeifer,F. (2004) GvpE- and GvpD-mediated transcription regulation of the p-gvp genes encoding gas vesicles in *Halobacterium salinarum*. *Microbiology* **150**: 1829-1838.
- Horne,M., Englert,C., Wimmer,C., and Pfeifer,F. (1991) A DNA region of 9 kbp contains all genes necessary for gas vesicle synthesis in halophilic archaeobacteria. *Mol Microbiol* **5**: 1159-1174.
- Huber,H., Hohn,M.J., Rachel,R., Fuchs,T., Wimmer,V.C., and Stetter,K.O. (2002) A new phylum of Archaea represented by a nanosized hyperthermophilic symbiont. *Nature* **417**: 63-67.
- Huber,H., Hohn,M.J., Stetter,K.O., and Rachel,R. (2003) The phylum Nanoarchaeota: present knowledge and future perspectives of a unique form of life. *Res Microbiol* **154**: 165-171.
- Inui,M., Vertes,A.A., and Yukawa,H. (1996) Reverse catabolite repression and the regulation of CO₂ fixation in *Rhodobacter* and related bacteria. *Research in Microbiology* **147**: 562-566.
- Jacob,F., and Monod,J. (1996) Genetic regulatory mechanisms in the synthesis of proteins. *J Mol Biol* **3**: 318-356.
- Joly,N., Bohm,A., Boos,W., and Richet,E. (2004) MalK, the ATP-binding cassette component of the *Escherichia coli* maltodextrin transporter, inhibits the transcriptional activator malt by antagonizing inducer binding. *J Biol Chem* **279**: 33123-33130.
- Jonuscheit,M., Martusewitsch,E., Stedman,K.M., and Schleper,C. (2003) A reporter gene system for the hyperthermophilic archaeon *Sulfolobus solfataricus* based on a selectable and integrative shuttle vector. *Mol Microbiol* **48**: 1241-1252.
- Kandler,O., and Hippe,H. (1977) Lack of peptidoglycan in the cell walls of *Methanosarcina barkeri*. *Arch Microbiol* **113**: 57-60.
- Kessler,A., Sezonov,G., Guijarro,J.I., Desnoues,N., Rose,T., Delepierre,M. *et al.* (2006) A novel archaeal regulatory protein, Sta1, activates transcription from viral promoters. *Nucleic Acids Res* **34**: 4837-4845.
- Kessler,P.S., Blank,C., and Leigh,J.A. (1998) The nif gene operon of the methanogenic archaeon *Methanococcus maripaludis*. *J Bacteriol* **180**: 1504-1511.
- Koning,S.M., Albers,S.V., Konings,W.N., and Driessen,A.J. (2002) Sugar transport in (hyper)thermophilic archaea. *Res Microbiol* **153**: 61-67.
- Koonin,E.V., Makarova,K.S., and Aravind,L. (2001) Horizontal gene transfer in prokaryotes: quantification and classification. *Annu Rev Microbiol* **55**: 709-742.

- Kosa,P.F., Ghosh,G., Decker,B.S., and Sigler,P.B. (1997) The 2.1-Å crystal structure of an archaeal preinitiation complex: TATA-box-binding protein/transcription factor (II)B core/TATA-box. *Proc Natl Acad Sci U S A* **94**: 6042-6047.
- Kruger,K., Hermann,T., Armbruster,V., and Pfeifer,F. (1998) The transcriptional activator GvpE for the halobacterial gas vesicle genes resembles a basic region leucine-zipper regulatory protein. *J Mol Biol* **279**: 761-771.
- Kuhnau,S., Reyes,M., Sievertsen,A., Shuman,H.A., and Boos,W. (1991) The activities of the Escherichia coli MalK protein in maltose transport, regulation, and inducer exclusion can be separated by mutations. *J Bacteriol* **173**: 2180-2186.
- Lange,U., and Hausner,W. (2004) Transcriptional fidelity and proofreading in Archaea and implications for the mechanism of TFS-induced RNA cleavage. *Mol Microbiol* **52**: 1133-1143.
- Langer,D., Hain,J., Thuriaux,P., and Zillig,W. (1995) Transcription in archaea: similarity to that in eucarya. *Proc Natl Acad Sci U S A* **92**: 5768-5772.
- Langer,D., Lottspeich,F., and Zillig,W. (1994) A subunit of an archaeal DNA-dependent RNA polymerase contains the S1 motif. *Nucleic Acids Res* **22**: 694.
- Langer,D., and Zillig,W. (1993) Putative tfIIs gene of Sulfolobus acidocaldarius encoding an archaeal transcription elongation factor is situated directly downstream of the gene for a small subunit of DNA-dependent RNA polymerase. *Nucleic Acids Res* **21**: 2251.
- Lee,H.S., Shockley,K.R., Schut,G.J., Conners,S.B., Montero,C.I., Johnson,M.R. *et al.* (2006) Transcriptional and biochemical analysis of starch metabolism in the hyperthermophilic archaeon Pyrococcus furiosus. *J Bacteriol* **188**: 2115-2125.
- Lee,S.J., Engelmann,A., Horlacher,R., Qu,Q., Vierke,G., Hebbeln,C. *et al.* (2003) TrmB, a sugar-specific transcriptional regulator of the trehalose/maltose ABC transporter from the hyperthermophilic archaeon Thermococcus litoralis. *J Biol Chem* **278**: 983-990.
- Lee,S.J., Moulakakis,C., Koning,S.M., Hausner,W., Thomm,M., and Boos,W. (2005) TrmB, a sugar sensing regulator of ABC transporter genes in Pyrococcus furiosus exhibits dual promoter specificity and is controlled by different inducers. *Mol Microbiol* **57**: 1797-1807.
- Leong,D., Boyer,H., and Betlach,M. (1988) Transcription of genes involved in bacterio-opsin gene expression in mutants of a halophilic archaeobacterium. *J Bacteriol* **170**: 4910-4915.
- Lie,T.J., Dodsworth,J.A., Nickle,D.C., and Leigh,J.A. (2007) Diverse homologues of the archaeal repressor NrpR function similarly in nitrogen regulation. *FEMS Microbiol Lett* **271**: 281-288.
- Lie,T.J., Wood,G.E., and Leigh,J.A. (2005) Regulation of nif expression in Methanococcus maripaludis: roles of the euryarchaeal repressor NrpR, 2-oxoglutarate, and two operators. *J Biol Chem* **280**: 5236-5241.
- Littlefield,O., Korkhin,Y., and Sigler,P.B. (1999) The structural basis for the oriented assembly of a TBP/TFB/promoter complex. *Proc Natl Acad Sci U S A* **96**: 13668-13673.
- Liu,W., Vierke,G., Wenke,A.K., Thomm,M., and Ladenstein,R. (2007) Crystal structure of the archaeal heat shock regulator from Pyrococcus furiosus: a molecular chimera representing eukaryal and bacterial features. *J Mol Biol* **369**: 474-488.

- Lubelska, J.M., Jonuscheit, M., Schleper, C., Albers, S.V., and Driessen, A.J. (2006) Regulation of expression of the arabinose and glucose transporter genes in the thermophilic archaeon *Sulfolobus solfataricus*. *Extremophiles* **10**(5); 383-91
- Marr, M.T., and Roberts, J.W. (2000) Function of transcription cleavage factors GreA and GreB at a regulatory pause site. *Mol Cell* **6**: 1275-1285.
- Martusewitsch, E., Sensen, C.W., and Schleper, C. (2000) High spontaneous mutation rate in the hyperthermophilic archaeon *Sulfolobus solfataricus* is mediated by transposable elements. *J Bacteriol* **182**: 2574-2581.
- Muller, B., Allmansberger, R., and Klein, A. (1985) Termination of a transcription unit comprising highly expressed genes in the archaeobacterium *Methanococcus voltae*. *Nucleic Acids Res* **13**: 6439-6445.
- Muller, S., and Klein, A. (2001) Coordinate positive regulation of genes encoding [NiFe] hydrogenases in *Methanococcus voltae*. *Mol Genet Genomics* **265**: 1069-1075.
- Napoli, A., Van der, O.J., Sensen, C.W., Charlebois, R.L., Rossi, M., and Ciaramella, M. (1999) An Lrp-like protein of the hyperthermophilic archaeon *Sulfolobus solfataricus* which binds to its own promoter. *J Bacteriol* **181**: 1474-1480.
- Newman, E.B., and Lin, R. (1995) Leucine-responsive regulatory protein: a global regulator of gene expression in *E. coli*. *Annu Rev Microbiol* **49**: 747-775.
- Nikolov, D.B., Chen, H., Halay, E.D., Usheva, A.A., Hisatake, K., Lee, D.K. *et al.* (1995) Crystal structure of a TFIIB-TBP-TATA-element ternary complex. *Nature* **377**: 119-128.
- O'Gara, F., Birkenhead, K., Boesten, B., and Fitzmaurice, A.M. (1989) Carbon metabolism and catabolite repression in *Rhizobium* spp. *FEMS Microbiol Rev* **5**: 93-101.
- Ouhammouch, M., Dewhurst, R.E., Hausner, W., Thomm, M., and Geiduschek, E.P. (2003) Activation of archaeal transcription by recruitment of the TATA-binding protein. *Proc Natl Acad Sci U S A* **100**: 5097-5102.
- Ouhammouch, M., and Geiduschek, E.P. (2005) An expanding family of archaeal transcriptional activators. *Proc Natl Acad Sci U S A* **102**: 15423-15428.
- Ouhammouch, M., Langham, G.E., Hausner, W., Simpson, A.J., El-Sayed, N.M., and Geiduschek, E.P. (2005) Promoter architecture and response to a positive regulator of archaeal transcription. *Mol Microbiol* **56**: 625-637.
- Palmer, J.R., and Daniels, C.J. (1995) In vivo definition of an archaeal promoter. *J Bacteriol* **177**: 1844-1849.
- Panagiotidis, C.H., Boos, W., and Shuman, H.A. (1998) The ATP-binding cassette subunit of the maltose transporter MalK antagonizes MalT, the activator of the *Escherichia coli* mal regulon. *Mol Microbiol* **30**: 535-546.
- Parvin, J.D., and Sharp, P.A. (1993) DNA topology and a minimal set of basal factors for transcription by RNA polymerase II. *Cell* **73**: 533-540.

- Parvin, J.D., Shykind, B.M., Meyers, R.E., Kim, J., and Sharp, P.A. (1994) Multiple sets of basal factors initiate transcription by RNA polymerase II. *J Biol Chem* **269**: 18414-18421.
- Peeters, E., Thia-Toong, T.L., Gigot, D., Maes, D., and Charlier, D. (2004) Ss-LrpB, a novel Lrp-like regulator of *Sulfolobus solfataricus* P2, binds cooperatively to three conserved targets in its own control region. *Mol Microbiol* **54**: 321-336.
- Peeters, E., Willaert, R., Maes, D., and Charlier, D. (2006) Ss-LrpB from *Sulfolobus solfataricus* condenses about 100 base pairs of its own operator DNA into globular nucleoprotein complexes. *J Biol Chem* **281**: 11721-11728.
- Pretz, M.G., van der, D.C., Albers, S.V., Schuurman-Wolters, G., and Driessen, A.J. (2008) Structural stability of GlcV, the nucleotide binding domain of the glucose ABC transporter of *Sulfolobus solfataricus*. *Biochim Biophys Acta* **1778**: 324-333.
- Qureshi, S.A., and Jackson, S.P. (1998) Sequence-specific DNA binding by the *S. shibatae* TFIIB homolog, TFB, and its effect on promoter strength. *Mol Cell* **1**: 389-400.
- Reines, D., Ghanouni, P., Gu, W., Mote, J., Jr., and Powell, W. (1993) Transcription elongation by RNA polymerase II: mechanism of SII activation. *Cell Mol Biol Res* **39**: 331-338.
- Reiter, W.D., Hudepohl, U., and Zillig, W. (1990) Mutational analysis of an archaeobacterial promoter: essential role of a TATA box for transcription efficiency and start-site selection in vitro. *Proc Natl Acad Sci U S A* **87**: 9509-9513.
- Roder, R., and Pfeifer, F. (1996) Influence of salt on the transcription of the gas-vesicle genes of *Haloferax mediterranei* and identification of the endogenous transcriptional activator gene. *Microbiology* **142 (Pt 7)**: 1715-1723.
- Rowlands, T., Baumann, P., and Jackson, S.P. (1994) The TATA-binding protein: a general transcription factor in eukaryotes and archaeobacteria. *Science* **264**: 1326-1329.
- Santangelo, T.J., Cubonova, L., James, C.L., and Reeve, J.N. (2007) TFB1 or TFB2 is sufficient for *Thermococcus kodakaraensis* viability and for basal transcription in vitro. *J Mol Biol* **367**: 344-357.
- Santangelo, T.J., and Reeve, J.N. (2006) Archaeal RNA polymerase is sensitive to intrinsic termination directed by transcribed and remote sequences. *J Mol Biol* **355**: 196-210.
- Santangelo, T.J., and Roberts, J.W. (2004) Forward translocation is the natural pathway of RNA release at an intrinsic terminator. *Mol Cell* **14**: 117-126.
- Schelert, J., Dixit, V., Hoang, V., Simbahan, J., Drozda, M., and Blum, P. (2004) Occurrence and characterization of mercury resistance in the hyperthermophilic archaeon *Sulfolobus solfataricus* by use of gene disruption. *J Bacteriol* **186**: 427-437.
- Schneider, E., and Hunke, S. (1998) ATP-binding-cassette (ABC) transport systems: functional and structural aspects of the ATP-hydrolyzing subunits/domains. *FEMS Microbiol Rev* **22**: 1-20.
- Schreiber, V., and Richet, E. (1999) Self-association of the *Escherichia coli* transcription activator MalT in the presence of maltotriose and ATP. *J Biol Chem* **274**: 33220-33226.

- Schut,G.J., Brehm,S.D., Datta,S., and Adams,M.W. (2003) Whole-genome DNA microarray analysis of a hyperthermophile and an archaeon: *Pyrococcus furiosus* grown on carbohydrates or peptides. *J Bacteriol* **185**: 3935-3947.
- Shibata,R., Bessho,Y., Shinkai,A., Nishimoto,M., Fusatomi,E., Terada,T. *et al.* (2007) Crystal structure and RNA-binding analysis of the archaeal transcription factor NusA. *Biochem Biophys Res Commun* **355**: 122-128.
- Shuman,H.A. (1982) Active transport of maltose in *Escherichia coli* K12. Role of the periplasmic maltose-binding protein and evidence for a substrate recognition site in the cytoplasmic membrane. *J Biol Chem* **257**: 5455-5461.
- Soppa,J. (1999b) Transcription initiation in Archaea: facts, factors and future aspects. *Mol Microbiol* **31**: 1295-1305.
- Soppa,J. (1999a) Normalized nucleotide frequencies allow the definition of archaeal promoter elements for different archaeal groups and reveal base-specific TFB contacts upstream of the TATA box. *Mol Microbiol* **31**: 1589-1592.
- Stiller,J.W., and Hall,B.D. (2002) Evolution of the RNA polymerase II C-terminal domain. *Proc Natl Acad Sci U S A* **99**: 6091-6096.
- Sun,J., and Klein,A. (2004) A lysR-type regulator is involved in the negative regulation of genes encoding selenium-free hydrogenases in the archaeon *Methanococcus voltae*. *Mol Microbiol* **52**: 563-571.
- Szabo,Z., Sani,M., Groeneveld,M., Zolghadr,B., Schelert,J., Albers,S.V. *et al.* (2007) Flagellar motility and structure in the hyperthermoacidophilic archaeon *Sulfolobus solfataricus*. *J Bacteriol* **189**: 4305-4309.
- Thomas,M.J., Platas,A.A., and Hawley,D.K. (1998) Transcriptional fidelity and proofreading by RNA polymerase II. *Cell* **93**: 627-637.
- Thomm,M. (1996) Archaeal transcription factors and their role in transcription initiation. *FEMS Microbiol Rev* **18**: 159-171.
- Thompson,D.K., Palmer,J.R., and Daniels,C.J. (1999) Expression and heat-responsive regulation of a TFIIB homologue from the archaeon *Haloferax volcanii*. *Mol Microbiol* **33**: 1081-1092.
- Tyree,C.M., George,C.P., Lira-DeVito,L.M., Wampler,S.L., Dahmus,M.E., Zawel,L., and Kadonaga,J.T. (1993) Identification of a minimal set of proteins that is sufficient for accurate initiation of transcription by RNA polymerase II. *Genes Dev* **7**: 1254-1265.
- Van de Vossenberg, Driessen,A.J., and Konings,W.N. (1998) The essence of being extremophilic: the role of the unique archaeal membrane lipids. *Extremophiles* **2**: 163-170.
- Van de Vossenberg, Ubbink-Kok,T., Elferink,M.G., Driessen,A.J., and Konings,W.N. (1995) Ion permeability of the cytoplasmic membrane limits the maximum growth temperature of bacteria and archaea. *Mol Microbiol* **18**: 925-932.
- van Hijum,S.A., de,J.A., Baerends,R.J., Karsens,H.A., Kramer,N.E., Larsen,R. *et al.* (2005) A generally applicable validation scheme for the assessment of factors involved in reproducibility and quality of DNA-microarray data. *BMC Genomics* **6**: 77.

- van Hijum, S.A., Garcia de la, N.J., Trelles, O., Kok, J., and Kuipers, O.P. (2003) MicroPreP: a cDNA microarray data pre-processing framework. *Appl Bioinformatics* **2**: 241-244.
- Verdon, G., Albers, S.V., Dijkstra, B.W., Driessen, A.J., and Thunnissen, A.M. (2003) Crystal structures of the ATPase subunit of the glucose ABC transporter from *Sulfolobus solfataricus*: nucleotide-free and nucleotide-bound conformations. *J Mol Biol* **330**: 343-358.
- Verdon, G., Albers, S.V., Dijkstra, B.W., Driessen, A.J., and Thunnissen, A.M. (2002) Purification, crystallization and preliminary X-ray diffraction analysis of an archaeal ABC-ATPase. *Acta Crystallogr D Biol Crystallogr* **58**: 362-365.
- Vierke, G., Engelmann, A., Hebbeln, C., and Thomm, M. (2003) A novel archaeal transcriptional regulator of heat shock response. *J Biol Chem* **278**: 18-26.
- Voorhorst, W.G., Gueguen, Y., Geerling, A.C., Schut, G., Dahlke, I., Thomm, M. *et al.* (1999) Transcriptional regulation in the hyperthermophilic archaeon *Pyrococcus furiosus*: coordinated expression of divergently oriented genes in response to beta-linked glucose polymers. *J Bacteriol* **181**: 3777-3783.
- Wettach, J., Gohl, H.P., Tschochner, H., and Thomm, M. (1995) Functional interaction of yeast and human TATA-binding proteins with an archaeal RNA polymerase and promoter. *Proc Natl Acad Sci U S A* **92**: 472-476.
- Wind, M., and Reines, D. (2000) Transcription elongation factor SII. *Bioessays* **22**: 327-336.
- Woese, C.R., and Fox, G.E. (1977) Phylogenetic structure of the prokaryotic domain: the primary kingdoms. *Proc Natl Acad Sci U S A* **74**: 5088-5090.
- Woese, C.R., Kandler, O., and Wheelis, M.L. (1990) Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. *Proc Natl Acad Sci U S A* **87**: 4576-4579.
- Worbs, M., Bourenkov, G.P., Bartunik, H.D., Huber, R., and Wahl, M.C. (2001) An extended RNA binding surface through arrayed S1 and KH domains in transcription factor NusA. *Mol Cell* **7**: 1177-1189.
- Worthington, P., Hoang, V., Perez-Pomares, F., and Blum, P. (2003) Targeted disruption of the alpha-amylase gene in the hyperthermophilic archaeon *Sulfolobus solfataricus*. *J Bacteriol* **185**: 482-488.
- Xie, Y., and Reeve, J.N. (2005) Regulation of tryptophan operon expression in the archaeon *Methanothermobacter thermoautotrophicus*. *J Bacteriol* **187**: 6419-6429.
- Zawel, L., Kumar, K.P., and Reinberg, D. (1995) Recycling of the general transcription factors during RNA polymerase II transcription. *Genes Dev* **9**: 1479-1490.
- Zillig, W., Stetter, K.O., Wunderl, S., Schulz, W., Priess, H., and Scholz, I. (1980) The *Sulfolobus*-*Caldariella* Group - Taxonomy on the Basis of the Structure of Dna-Dependent Rna-Polymerases. *Archives of Microbiology* **125**: 259-269.
- Zimmermann, P., and Pfeifer, F. (2003) Regulation of the expression of gas vesicle genes in *Haloferax mediterranei*: interaction of the two regulatory proteins GvpD and GvpE. *Mol Microbiol* **49**: 783-794.

References

Zolghadr,B., Weber,S., Szabo,Z., Driessen,A.J., and Albers,S.V. (2007) Identification of a system required for the functional surface localization of sugar binding proteins with class III signal peptides in *Sulfolobus solfataricus*. *Mol Microbiol* **64**: 795-806.